

# Seralutinib demonstrated *in vitro* reduction of vascular inflammatory drivers underlying pulmonary hypertension


Zhaoqing Ding<sup>1</sup>, Elisa Schiavi<sup>2</sup>, Silvia Cantoni<sup>2</sup>, Ravikumar Sitapara<sup>1</sup>, Eduardo Garcia<sup>1</sup>, Giulia Nuozzi<sup>2</sup>, Jessica Burroughs-Garcia<sup>2</sup>, Robin Osterhout<sup>1</sup>, Lawrence S. Zisman<sup>1</sup>, Sidra Hoffman<sup>1</sup>, Richard Aranda<sup>1</sup>, Daniela Miglietta<sup>2</sup>, Stephan Rosenkranz<sup>3</sup>, Anna R. Hemnes<sup>4</sup>, Adam J. Byrne<sup>5</sup>, Jean-Marie Bruey<sup>1</sup>, Cormac McCarthy<sup>6</sup>

<sup>1</sup>Gossamer Bio, Inc., San Diego, CA, USA; <sup>2</sup>Chiesi Farmaceutici S.p.A., Parma, Italy; <sup>3</sup>Department of Cardiology - Internal Medicine III, Heart Center, University Hospital Cologne, Cologne, Germany; Cologne Cardiovascular Research Center (CCRC), University of Cologne, Cologne, Germany; <sup>4</sup>Vanderbilt University, Vanderbilt University Medical Center, Nashville, TN, USA; <sup>5</sup>University College Dublin, School of Medicine, Dublin, Ireland; <sup>6</sup>University College Dublin, School of Medicine, St. Vincent's Hospital, Dublin, Ireland

## BACKGROUND


- Inflammation and immune dysregulation contribute to vascular remodeling in group 1 and 3 pulmonary hypertension subtypes<sup>1,2</sup>
- The degree of perivascular immune infiltration correlates with vascular remodeling<sup>2-5</sup>
- Cytokines and chemokines produced by infiltrating immune cells contribute to endothelial injury, smooth muscle cell proliferation, and fibrogenesis/fibroblast activation<sup>6-8</sup>
- Seralutinib, a potent inhaled PDGFR, CSF1R, and c-KIT tyrosine kinase inhibitor (**Figure 1**), significantly reduced pulmonary vascular resistance in the phase 2 TORREY study in pulmonary arterial hypertension (PAH; NCT04456998)<sup>9</sup>
- Seralutinib treatment was associated with an anti-inflammatory proteomic signature after 24 weeks of treatment in TORREY<sup>10</sup> and in the SU5416/Hypoxia model of PAH<sup>11</sup>
- This study assessed seralutinib's impact on inflammation in primary human cellular immune assays

**Figure 1. Seralutinib mechanism of action**




## RESULTS

**Figure 2. Seralutinib reduced production of inflammatory cytokines and chemokines in *in vitro* models of primary human vascular inflammation**



**Figure 3. Seralutinib reduced TNF-α, IL-8, and MCP-1 production from activated primary human innate and adaptive immune cells**



Freshly isolated human PBMCs (n=4) were stimulated with either LPS (innate) or anti-CD3/anti-CD28 (adaptive; TCR) and treated with seralutinib at noted concentrations. MCP-1, IL-8, and TNF-α were measured from supernatant at 18 h (LPS) or 48 h (TCR). Percent inhibition was calculated compared to average of vehicle control per analyte measured. Data are expressed as mean ± SEM (n=4). Statistical analysis was performed using one-way ANOVA with Dunnett's multiple comparisons test. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001, and \*\*\*\*p<0.0001 versus stimulation control. ANOVA, analysis of variance; IL-8, interleukin 8; MCP-1, monocyte chemoattractant protein 1; PBMC, peripheral blood mononuclear cell; SEM, standard error of the mean; TCR, T-cell receptor; TNF-α, tumor necrosis factor α.

## METHODS

### *In vitro* system of primary human vascular inflammation

- Healthy primary human cell-based BioMAP® systems were used to assess immune and tissue-specific responses. Cells from 2-6 donors were pooled for each assay
  - LPS System: Peripheral blood mononuclear cells (PBMCs) co-cultured with human umbilical vein endothelial cells (HUVECs), stimulated with lipopolysaccharide, LPS (a TLR4 ligand) for 24 h
  - SAg System: PBMCs co-cultured with HUVECs, stimulated with super antigen, SAg (a T-cell receptor [TCR] cross-linker) for 96 h
- Soluble biomarkers were quantified using Homogeneous Time Resolved Fluorescence (HTRF®)

### *In vitro* system of primary human immune cell activation

- Freshly isolated PBMCs from 4 healthy donors were pretreated with seralutinib for 1 h and stimulated with:
  - LPS from *E. coli* O111:B4 (Invivogen) for 18 h (innate immune activation)
  - T Cell TransAct™ (Miltenyi Biotec) (activation via CD3 and CD28) for 48 h (adaptive immune activation)
- Soluble markers were measured by Luminex® assays (R&D Systems)

## CONCLUSIONS

- Seralutinib demonstrated anti-inflammatory effects across different human models of *in vitro* vascular inflammation and innate and adaptive immune activation
- In combination with clinical biomarker data, these findings highlight the potential of seralutinib to reduce the hyperactive vascular inflammation underlying pulmonary hypertension
- Further investigation of these mechanisms is planned

**References:** 1 Mocumbi A, et al. *Nat Rev Dis Primers*. 2024;10(1):1. 2 Liu SF, et al. *Front Immunol*. 2022;13:959209. 3 Dorfmueller P, et al. *Eur Respir J*. 2003;22(2):358-363. 4 Stacher E, et al. *Am J Respir Crit Care Med*. 2012;186(3):261-272. 5 Ferrian S, et al. *Am J Respir Crit Care Med*. 2024;209(2):206-218. 6 Bell RD, et al. *Arthritis Rheumatol*. 2020;72(10):1759-1770. 7 Sanchez O, et al. *Am J Respir Crit Care Med*. 2007;176(10):1041-1047. 8 Soon E, et al. *Circulation*. 2010;122(9):920-927. 9 Frantz RP, et al. *Lancet Respir Med*. 2024;12(7):523-534. 10 Ghofrani HA, et al. *Am J Respir Crit Care Med*. 2024;209:A7383. 11 Galkin A, et al. *Eur Respir J*. 2022;60(6):2102356.

**Disclosures:** ZD is an employee of, and owns stocks or stock options from, Gossamer Bio, Inc.  
**Research supported by:** Chiesi Farmaceutici S.p.A. and Gossamer Bio, Inc.



The ERS is not responsible for and does not endorse the data and information presented on external sites.