

Mapping the Diagnostic Journey of Patients With Pulmonary Hypertension Associated With Interstitial Lung Disease (PH-ILD): A Retrospective U.S. Claims Analysis

Kristin B. Highland¹, Sandy Nguyen², Jessica Schwartz³,
Samantha Rothstein³, Nidhishant Dixit³, Joyce Louie³, Nicole
Mayer³, Bara' Momani³, Lawrence Ho²

¹Cleveland Clinic Lerner College of Medicine, Cleveland, OH, USA; ²Gossamer Bio, Inc., San Diego, CA, USA; ³MMIT, Norstella Consulting & Analytics, London, UK

Research supported by: Chiesi Farmaceutici S.p.A. and Gossamer Bio, Inc.

**PHenomenal
Hope 2025**

Knowledge, Research & Advocacy in PH

**Boston, MA
December 5, 2025**

Background



- PH-ILD is a severe, progressive, and life-limiting cardiopulmonary condition characterized by elevated pulmonary artery pressure in concert with interstitial inflammation and fibrosis¹
- Timely diagnosis is difficult and often delayed owing to nonspecific and frequently overlapping symptoms between ILD and PH, a low index of suspicion among non-PH specialists, and lack of standardized diagnostic algorithms for WHO Group 3 PH^{1,2}
- This study characterizes the diagnostic journey of PH-ILD patients using real-world claims data, focusing on the diagnosis of PH in patients with ILD

Methods



- A retrospective cohort study was conducted using the NorstellLinQ fully integrated claims and electronic health record dataset from January 2019 to December 2024

Inclusion Criteria


- Adults with ≥ 1 claim for ILD and ≥ 1 claim for PH
- PH-ILD was defined using ICD-10 codes indicative of both ILD and PH, excluding other PH groups

Exclusion Criteria

- COPD or emphysema
- CV conditions associated with WHO Group 2 PH
- Heart failure in medical record prior to PH diagnosis

- Time intervals were calculated from symptom presentation to PH diagnosis
- The diagnosing physician's specialty, as well as diagnostic procedures, were examined


Patient cohort selection



Patient demographics




- Among 18,666 PH-ILD patients identified, the majority were female, White, and age ≥ 65 years



There was an increase in female predominance at each step of the patient cohort funnel



Gender by patient cohort



Within the claims analysis cohort, a large proportion were classified as having CTD-ILD (33%)



ILD Subtype	Patients, n (%) [*]
Connective tissue disease	6,218 (33)
IPF	1,881 (10)
Toxin	1,112 (6)
Hypersensitivity pneumonitis	304 (2)
Occupational / environmental	164 (1)
Radiation	76 (0.4)
Other ILDs ^a	10,118 (54)

Expected patterns of ILD demographics




- IPF accounts for ~ 1/3 of ILD cases and is strongly male-predominant, which has historically shaped perceptions of ILD demographics¹
- However, most other ILD subtypes—including CTD-ILD, IPAF, NSIP, and fibrotic hypersensitivity pneumonitis—skew female due to their autoimmune associations^{2–8}
- Patients with emphysema, including CPFE, which exhibits a male predominance of 9:1,⁹ were excluded from this cohort
- These patterns may explain why this PH-ILD cohort was predominantly female, driven by the high proportion of non-IPF ILDs

1. Maher TM. *JAMA*. 2024;331(19):1655-1665. 2. Gezgin T, et al. *Eur Research J*. 2023;9(6):1360-1367. 3. Kaul B, et al. *Front Med (Lausanne)*. 2021;8:751181. 4. Fernandes L, et al. *Front Med (Lausanne)*. 2019;6:209. 5. Sebastiani M, et al. *Biomedicines*. 2020;9(1):17. 6. Belloli EA, et al. *Respirology*. 2016;21(2):259-268. 7. Creamer AW, Barratt SL. *Eur Respir Rev*. 2020;29(156):190167. 8. Torres-Rojas MB, et al. *Pulm Circ*. 2025;15(1):e70035. 9. Jankowich MD, Rounds SIS. *Chest*. 2012;141(1):222-231.

PH diagnosis occurrence relative to ILD diagnosis




Most patients (65%) received a PH diagnosis at the same time as, or after, being diagnosed with ILD



Symptom occurrence in the years prior to and after PH diagnosis



The most common symptoms preceding PH diagnosis were dyspnea (20%), dizziness (14%), edema (14%), and palpitations (12%)



Specialties of HCPs who diagnosed PH




A substantial percentage of non-experts are diagnosing PH.

HCPs diagnosing PH-ILD	Patients, n (%)
Diagnosed by PH-focused specialties	10,189 (55)
Cardiology	5,371 (29)
Pulmonology	3,610 (19)
Rheumatology	1,070 (6)
Other	53 (<1)
Diagnosed by non-PH-focused specialties^a	8,477 (45)

^aFamily/internal medicine (21%), nurse practitioner-other (4%), hospitalist (3%), radiology-diagnostic (3%), critical care (3%), pediatrics (2%).
HCP, healthcare professional; ILD, interstitial lung disease; n, number; PH, pulmonary hypertension.

Most common diagnostic tests performed before or in the same year as PH diagnosis



- 22% of patients used supplemental oxygen before or in the same year as PH diagnosis

Pulmonologists most frequently ordered diagnostic tests or supplemental O₂ (except for echocardiography)



Diagnostic test or supportive measure administered by specialist type



Echocardiography

- Cardiologists 23%
- Pulmonologists 18%



CT scan

- Pulmonologists 21%
- Cardiologists 21%



Right heart catheterization

- Pulmonologists 29%
- Cardiologists 15%



Spirometry

- Pulmonologists 29%
- Cardiologists 21%



Supplemental oxygen use

- Pulmonologists 26%
- Cardiologists 18%



6-minute walk test

- Pulmonologists 34%
- Cardiologists 19%

Limitations



- Patients had ≥ 1 claim for PH and ≥ 1 claim for ILD but there was no requirement for confirmatory diagnosis claims for each condition
- Generic ILD codes may contribute to the lower-than-expected percentage of IPF patients in this cohort
- No physician chart review was performed to differentiate WHO Group 3 PH from WHO Group 1 PH (for example, in patients with CTD)
- Sarcoidosis was not explicitly excluded, and therefore some Group 5 PH patients may have been included in this cohort

Conclusions



- This analysis offers insights into symptom presentation, physician specialties, and modalities used for PH-ILD diagnosis
- The female predominance in this cohort may be explained by the high representation of CTD and other non-IPF ILD patients and the exclusion of CPFE patients in the analysis
- There was a delay of up to 1.6 years between initial symptom presentation and PH diagnosis, likely attributed to the overlapping symptomatology of PH and ILD
- A high percentage of diagnoses was made by non-PH specialists in acute and primary care settings, contributing to potential additional delays in referral to appropriate specialty care centers
- Key diagnostic modalities—including RHC, PFTs, and 6MWT—remain underutilized and may represent missed opportunities for comprehensive evaluation and timely intervention in this high-risk population
- Opportunities exist to accelerate disease awareness, recognition, and referral to reduce diagnostic delays and optimize outcomes for this high-risk population